

Integration of Dirac-Jacobi structures

David Iglesias Ponte* and Aïssa Wade†‡

Department of Mathematics, Penn State University

Abstract

We study precontact groupoids whose infinitesimal counterparts are Dirac-Jacobi structures. These geometric objects generalize contact groupoids. We also explain the relationship between precontact groupoids and homogeneous presymplectic groupoids. Finally, we present some examples of precontact groupoids.

1 Introduction

Presymplectic groupoids, introduced and studied in [2], are global counterparts of Dirac structures. They allow to extend the well-known correspondence between symplectic groupoids and Poisson manifolds to the context of Dirac geometry. Moreover, they provide a framework for a unified formulation of various notions of momentum maps ([1]). On the other hand, Dirac-Jacobi structures (called $\mathcal{E}^1(M)$ -Dirac structures in [11]) include both Dirac and Jacobi structures. They naturally appeared in the geometric prequantization of Dirac manifolds [13, 14].

In this paper, our aim is to investigate the integrability problem for Dirac-Jacobi structures. This work is motivated by the fact that many Dirac manifolds can be quantized through their integrating Lie groupoids. We show that the global counterparts of Dirac-Jacobi manifolds are what we call here precontact groupoids. In particular, we recover the integrability of Jacobi structures [4]. We also prove that there is a one-to-one correspondence between precontact groupoids and homogeneous presymplectic groupoids. Moreover, the precontact groupoid $\tilde{G}$ associated with an integrable Dirac structure L_0 on M is just the prequantization of the presymplectic groupoid G associated with L_0 (that is, the central extension of Lie groupoids $M \times S^1 \rightarrow \tilde{G} \rightarrow G$ satisfying some compatibility conditions), provided that the canonical Dirac Jacobi structure L on M corresponding L_0 is integrable, see Section 5.2. We should mention that M. Zambon and C. Zhu independently study the geometry of prequantization spaces [14].

*e-mail: iglesias@math.psu.edu

†e-mail: wade@math.psu.edu

‡The first author is partially supported by MCYT grant BFM2003-01319

Here is an outline of the paper. In Sections 2 and 3, we give some definitions and results needed to establish our results. Section 4 contains our main results (Theorems 4.2 and 4.3). In Section 5, we give some examples of precontact groupoids.

2 Basic definitions and results

2.1 Dirac structures and presymplectic groupoids

Let M be a smooth n -dimensional manifold. There is a natural symmetric pairing $\langle \cdot, \cdot \rangle$ on the vector bundle $TM \oplus T^*M$ given by

$$\langle X_1 + \xi_1, X_2 + \xi_2 \rangle = \frac{1}{2} \left(\xi_1(X_2) + \xi_2(X_1) \right).$$

Furthermore, the space of smooth sections of $TM \oplus T^*M$ is endowed with the Courant bracket, which is defined by

$$[X_1 + \xi_1, X_2 + \xi_2] = [X_1, X_2] + \mathcal{L}_{X_1}\xi_2 - i_{X_2}d\xi_1,$$

for any $X_1 + \xi_1, X_2 + \xi_2 \in \Gamma(TM \oplus T^*M)$.

Definition 2.1 [2, 5] A **Dirac structure** on a smooth manifold M is a subbundle L of $TM \oplus T^*M$ which is maximally isotropic with respect to the symmetric pairing $\langle \cdot, \cdot \rangle$ and whose space of sections is closed under the Courant bracket.

Let L_M and L_N be Dirac structures on M and N , respectively. We say that a smooth map $F : M \rightarrow N$ is a (forward) **Dirac map** if $L_N = F_*(L_M)$, where

$$F_*(L_M) = \{ (dF)(X) + \xi \mid X + F^*\xi \in L_M \}.$$

Recall that any Dirac structure L has an induced Lie algebroid structure: the Lie bracket on $\Gamma(L)$ is just the restriction of the Courant bracket and the anchor map is the restriction of the first projection to L , i.e. $pr_1|_L : L \rightarrow TM$. Now, suppose that L is a Dirac structure which is isomorphic to the Lie algebroid of a Lie groupoid G . Such a Lie groupoid is called an **integration** of the Lie algebroid L . Then, there exists an induced closed 2-form on G with some additional properties. More precisely,

Definition 2.2 [2] A **presymplectic groupoid** is a pair (G, ω) which consists of a groupoid $G \overset{\alpha}{\rightrightarrows} M$ such that $\dim(G) = 2 \dim(M)$, and a 2-form $\omega \in \Omega^2(G)$ satisfying the following conditions:

- (i) ω is closed, i.e. $d\omega = 0$.
- (ii) ω is multiplicative, that is, $m^*\omega = pr_1^*\omega + pr_2^*\omega$.
- (iii) $\text{Ker}(\omega_x) \cap \text{Ker}(d\alpha)_x \cap \text{Ker}(d\beta)_x = \{0\}$, for all $x \in M$.

We say that the presymplectic groupoid (G, ω) is **homogeneous** if there exists a multiplicative vector field Z such that $\mathcal{L}_Z \omega = \omega$.

The relationship between Dirac structures and presymplectic groupoids is provided by the following result:

Proposition 2.3 [2] *Given a presymplectic groupoid (G, ω) , there is a canonical Dirac structure L on M which is isomorphic to the Lie algebroid AG of G , and such that the target map $\beta : (G, L_\omega) \rightarrow (M, L)$ is a Dirac map, while the source map $\alpha : (G, L_\omega) \rightarrow (M, L)$ is anti-Dirac.*

Conversely, suppose that L is a Dirac structure on M whose associated Lie algebroid is integrable, and let $G(L)$ be its α -simply connected integration. Then, there exists a unique 2-form ω such that $(G(L), \omega)$ is a presymplectic groupoid, the target map is a Dirac map, and the source map is anti-Dirac.

2.2 Dirac-Jacobi structures

Let M be a smooth n -dimensional manifold. There is a natural bilinear operation $\langle \cdot, \cdot \rangle$ on the vector bundle $\mathcal{E}^1(M) = (TM \times \mathbb{R}) \oplus (T^*M \times \mathbb{R})$ defined by:

$$\left\langle (X_1, f_1) + (\xi_1, g_1), (X_2, f_2) + (\xi_2, g_2) \right\rangle = \frac{1}{2}(i_{X_2}\xi_1 + i_{X_1}\xi_2 + f_1g_2 + f_2g_1),$$

for any $(X_\ell, f_\ell) + (\xi_\ell, g_\ell) \in \Gamma(\mathcal{E}^1(M))$, with $\ell = 1, 2$. In addition, the space of smooth sections of $\mathcal{E}^1(M)$ is equipped with an $\mathbb{R}$ -bilinear operation which can be viewed as an extension of the Courant bracket on $TM \oplus T^*M$, i.e.

$$\begin{aligned} [(X_1, f_1) + (\xi_1, g_1), (X_2, f_2) + (\xi_2, g_2)] &= \left([X_1, X_2], X_1(f_2) - X_2(f_1) \right) \\ &\quad + \left(\mathcal{L}_{X_1}\xi_2 - \mathcal{L}_{X_2}\xi_1 + \frac{1}{2}d(i_{X_2}\xi_1 - i_{X_1}\xi_2) \right. \\ &\quad \left. + f_1\xi_2 - f_2\xi_1 + \frac{1}{2}(g_2df_1 - g_1df_2 - f_1dg_2 + f_2dg_1), \right. \\ &\quad \left. X_1(g_2) - X_2(g_1) + \frac{1}{2}(i_{X_2}\xi_1 - i_{X_1}\xi_2 - f_2g_1 + f_1g_2) \right), \end{aligned}$$

for any $(X_\ell, f_\ell) + (\xi_\ell, g_\ell) \in \Gamma(\mathcal{E}^1(M))$ with $\ell = 1, 2$. For an alternative description of this bracket, see [9].

Definition 2.4 [11] *A **Dirac-Jacobi structure** is a subbundle L of $\mathcal{E}^1(M)$ which is maximally isotropic with respect to $\langle \cdot, \cdot \rangle$ and such that $\Gamma(L)$ is closed under the extended Courant bracket $[\cdot, \cdot]$.*

*Let L_M (resp., L_N) be a Dirac-Jacobi structure on M (resp., N). We say that a smooth surjective map $F : M \rightarrow N$ is a (forward) **Dirac-Jacobi map** if $L_N = F_*(L_M)$, where*

$$F_*(L_M) = \{((dF)(X), f) + (\xi, g) \mid (X, f \circ F) + (F^*\xi, g \circ F) \in L_M\}.$$

Basic examples of Dirac-Jacobi structures are Dirac and Jacobi structures on M (this explains the terminology introduced in [7]).

2.3 Action Lie algebroids and 1-cocycles

It is known that, given any a Lie algebroid $(A, \llbracket \cdot, \cdot \rrbracket, \rho)$ over M and any 1-cocycle $\phi \in \Gamma(A^*)$, there is an associated Lie algebroid over $M \times \mathbb{R}$, denoted by $(A \times_\phi \mathbb{R}, \llbracket \cdot, \cdot \rrbracket^\phi, \rho^\phi)$, where the smooth sections of $A \times_\phi \mathbb{R}$ are of the form $\bar{X}(x, t) = X_t(x)$, with $X_t \in \Gamma(A)$ for all $t \in \mathbb{R}$, and

$$\begin{aligned} \llbracket \bar{X}, \bar{Y} \rrbracket^\phi(x, t) &= \llbracket X_t, Y_t \rrbracket(x) + \phi(X_t)(x) \frac{\partial \bar{Y}}{\partial t} - \phi(Y_t)(x) \frac{\partial \bar{X}}{\partial t}, \\ \rho^\phi(\bar{X})(x, t) &= \rho(X_t)(x) + \phi(X_t)(x) \frac{\partial}{\partial t}, \end{aligned} \quad (1)$$

where $\frac{\partial \bar{X}}{\partial t} \in \Gamma(A \times_\phi \mathbb{R})$ denotes the derivative of $\bar{X}$ with respect to t .

Remark 2.5 If L is a Dirac-Jacobi structure then the restriction of the extended Courant bracket to sections of L together with the canonical projection of L onto TM make L into a Lie algebroid over M . In addition, $\phi \in \Gamma(L^*)$ defined by

$$\phi(v) = f, \text{ for } v = (X, f) + (\xi, g) \in \Gamma(L), \quad (2)$$

is a 1-cocycle for the Lie algebroid cohomology (see [8]). On the other hand, it is known that a Dirac-Jacobi structure L on M corresponds to a Dirac structure $\tilde{L}$ on $M \times \mathbb{R}$ given by

$$\tilde{L} = \left\{ \left(X + f \frac{\partial}{\partial t} \right) + \left(e^t(\xi + g dt) \right) \mid (X, f) + (\xi, g) \in L \right\}. \quad (3)$$

Moreover, $\tilde{L}$ is isomorphic to $L \times_\phi \mathbb{R}$.

2.4 Conformal classes of Dirac-Jacobi structures

Let L be a Dirac-Jacobi structure on M and let φ be a smooth nowhere vanishing function on M . We set $\mu = d \ln |\varphi|$. Consider the vector bundle L_φ over M whose space of smooth sections is given by

$$\Gamma(L_\varphi) = \{(X, f - \mu(X)) + \varphi(\xi + g\mu, g) \mid (X, f) + (\xi, g) \in \Gamma(L)\}.$$

One can easily check that L_φ is also a Dirac-Jacobi structure on M . The correspondence $(L, \varphi) \mapsto L_\varphi$ is called a conformal change. For any fixed L , the family of all (L_φ) is called a conformal class of Dirac-Jacobi structures. For instance, when L comes from a presymplectic form ω then L_φ is nothing but the Dirac-Jacobi structure associated with $(\varphi\omega, d \ln |\varphi|)$ (see [11, 12] for more details).

3 Precontact groupoids

Definition 3.1 Let $G \xrightleftharpoons[\beta]{\alpha} M$ be a Lie groupoid such that $\dim(G) = 2 \dim(M) + 1$. A **precontact groupoid structure** on G is given by a pair (η, σ) consisting of a 1-form η

and a multiplicative function σ (i.e., $\sigma(gh) = \sigma(g) + \sigma(h)$) such that

$$m^*\eta = pr_1^*\eta + pr_1^*(e^\sigma)pr_2^*\eta. \quad (4)$$

$$\text{Ker}(d\eta_x) \cap \text{Ker}(\eta_x) \cap \text{Ker}(d\alpha)_x \cap \text{Ker}(d\beta)_x = \{0\}, \text{ for all } x \in M. \quad (5)$$

Two precontact structures (η, σ) and (η', σ') on G are equivalent if there exists a nowhere vanishing function $\varphi : M \rightarrow \mathbb{R}$ such that

$$\eta' = (\varphi \circ \alpha) \eta, \quad \sigma' = \sigma + \ln \left| \frac{\varphi \circ \alpha}{\varphi \circ \beta} \right|$$

Now, consider a Lie groupoid $G \rightrightarrows M$ together with a multiplicative function σ . One can define a right action of G on the canonical projection $\pi_1 : M \times \mathbb{R} \rightarrow M$ as follows:

$$(x, t) \cdot g = (\alpha(g), t + \sigma(g)), \text{ for } (x, t, g) \in M \times \mathbb{R} \times G, \text{ such that } \beta(g) = x.$$

Therefore, we have the corresponding action groupoid $G \times \mathbb{R} \rightrightarrows M \times \mathbb{R}$, denoted by $G \times_\sigma \mathbb{R}$, with structural functions given by

$$\begin{aligned} \alpha_\sigma(g, t) &= (\alpha(g), \sigma(g) + t), \\ \beta_\sigma(h, s) &= (\beta(h), s), \\ m_\sigma((g, t), (h, s)) &= (gh, t), \text{ if } \alpha_\sigma(g, t) = \beta_\sigma(h, s). \end{aligned} \quad (6)$$

We denote by $(AG, \llbracket \cdot, \cdot \rrbracket, \rho)$ the Lie algebroid of G . The multiplicative function σ induces a 1-cocycle ϕ on AG given by

$$\phi(x)(X_x) = X_x(\sigma), \text{ for } x \in M \text{ and } X_x \in A_x G. \quad (7)$$

In addition, we can identify the Lie algebroid of the Lie groupoid $G \times_\sigma \mathbb{R}$ with $AG \times_\phi \mathbb{R}$. Conversely, one has the following

Proposition 3.2 [4] *Let L be a Lie algebroid over M , ϕ be a 1-cocycle and $L \times_\phi \mathbb{R}$ the Lie algebroid given by Equation (1). Then, L is integrable if and only if $L \times_\phi \mathbb{R}$ is integrable. Moreover, if $G(L)$ (resp., $G(L \times_\phi \mathbb{R})$) is the α -simply connected integration of L (resp., $L \times_\phi \mathbb{R}$) and σ is the multiplicative function associated with ϕ , then $G(L \times_\phi \mathbb{R}) \cong G(L) \times_\sigma \mathbb{R}$.*

There is a correspondence between precontact and presymplectic groupoids. Indeed, one has the following proposition:

Proposition 3.3 *Let $G \rightrightarrows M$ be a Lie groupoid and σ a multiplicative function on G . There is a one-to-one correspondence between precontact groupoids on (G, σ) and homogeneous presymplectic groupoids on $G \times_\sigma \mathbb{R}$.*

Proof: We know that there exists a one-to-one correspondence between 1-forms on a manifold M and 2-forms on $M \times \mathbb{R}$ homogeneous with respect to $\frac{\partial}{\partial t}$. More precisely, if η is a 1-form on M then $\omega = d(e^t \eta)$ is a homogeneous 2-form on $M \times \mathbb{R}$. Conversely, assume that ω is homogeneous and set $\eta = i_{\frac{\partial}{\partial t}} \omega$. One can check that $\mathcal{L}_{\frac{\partial}{\partial t}} \eta = \eta$. Hence η can be identified with a 1-form on M . Using the relation $\omega = d(e^t \eta)$, it is straightforward to prove that conditions (ii) and (iii) in Definition 2.2 are equivalent to Equations (4) and (5). $\blacksquare$

4 Integration of Dirac-Jacobi structures

In this section, we show that precontact groupoids are the global objects corresponding to Dirac-Jacobi structures. First, note that the following lemma which is an immediate consequence of Remark 2.5 and Proposition 3.2.

Lemma 4.1 *A Dirac-Jacobi structure L is integrable if and only if its associated Dirac structure $\tilde{L} \subset T(M \times \mathbb{R}) \oplus T^*(M \times \mathbb{R})$ is integrable.*

Theorem 4.2 *Let L be an integrable Dirac-Jacobi structure on M , and let $G(L)$ be its α -simply connected integration. There exists a multiplicative function σ and a 1-form η such that $(G(L), \eta, \sigma)$ is a precontact groupoid. Furthermore, any conformal class of integrable Dirac-Jacobi structures on M induces a conformal class of precontact groupoid structures.*

Proof: Suppose L is an integrable Dirac-Jacobi structure on M . We denote by $\phi \in \Gamma(L^*)$ the 1-cocycle defined by Equation (2). By integration, the 1-cocycle ϕ induces a multiplicative function σ on $G(L)$. Denote by $\tilde{L}$ the Dirac structure on $M \times \mathbb{R}$ associated with L given by Equation (3). Applying Proposition 2.3, one gets a presymplectic groupoid $(G(\tilde{L}), \omega)$. Moreover, one gets from Remark 2.5 and Proposition 3.2 that

$$G(\tilde{L}) \cong G(L \times_{\phi} \mathbb{R}) \cong G(L) \times_{\sigma} \mathbb{R}.$$

Observe that ω is homogeneous with respect to the multiplicative vector field $\frac{\partial}{\partial t}$. Indeed, the differentiable family of diffeomorphisms $\psi_s : M \times \mathbb{R} \rightarrow M \times \mathbb{R}$ defined by $\psi_s(x, t) = (x, s + t)$ consists of Dirac maps, i.e. $(\psi_s)_*(\tilde{L}) = \tilde{L}$. Furthermore, by integration and using the flow $\psi_s : G(L) \times \mathbb{R} \rightarrow G(L) \times \mathbb{R}$ $(g, t) \mapsto (g, t + s)$ of the vector field $\frac{\partial}{\partial t}$, we get

$$\mathcal{L}_{\frac{\partial}{\partial t}} \omega = \omega.$$

Thus, using Proposition 3.3, we deduce that there exists a 1-form η such that $(G(L), \eta, \sigma)$ is precontact. Recall that, for every nowhere vanishing function φ on M , there is an equivalent Dirac-Jacobi structure on M whose space of sections is given by

$$\Gamma(L_{\varphi}) = \{(X, f - \mu(X)) + \varphi(\xi + g\mu, g) \mid (X, f) + (\xi, g) \in \Gamma(L)\}.$$

Consider the 1-cocycle $\phi_\varphi \in \Gamma(L_\varphi^*)$ defined as follows

$$\phi_\varphi(e) = f - \mu(X), \quad \text{for all } e = (X, f - \mu(X)) + \varphi(\xi + g\mu, g).$$

We have a natural commutative diagram of vector bundle morphisms:

$$\begin{array}{ccc} L & \xrightarrow{\cong} & L_\varphi \\ \downarrow & & \downarrow \\ L \times_\phi \mathbb{R} & \xrightarrow{\cong} & L_\varphi \times_{\phi_\varphi} \mathbb{R} \end{array}$$

Moreover, L_φ induces a precontact structure $(\eta_\varphi, \sigma_\varphi)$ on $G(L_\varphi) \cong G(L)$ given by

$$\eta_\varphi = (\varphi \circ \alpha) \eta \quad \sigma_\varphi = \sigma + \ln \left| \frac{\varphi \circ \alpha}{\varphi \circ \beta} \right|.$$

Thus, any conformal class of integrable Dirac-Jacobi structures on M induces a conformal class of precontact groupoid structures. ■

Conversely, we have the following result.

Theorem 4.3 *Let (G, η, σ) is a precontact groupoid over M . Then, there exists a canonical Dirac-Jacobi structure L_M on M which is isomorphic to the Lie algebroid AG . Moreover, $\beta : G \rightarrow M$ is a Dirac-Jacobi map.*

Proof: By Proposition 3.3, one has the presymplectic groupoid $(G \times_\sigma \mathbb{R}, \omega = d(e^t \eta))$ over $M \times \mathbb{R}$. Then, using Proposition 2.3, one gets a Dirac structure $L_{M \times \mathbb{R}}$ on $M \times \mathbb{R}$ such that β_σ is a Dirac map. Thus,

$$\begin{aligned} L_{M \times \mathbb{R}} &= \left\{ d\beta_\sigma \left(X + F \frac{\partial}{\partial t} \right) + (\xi + Gdt) \mid i_{(X+F \frac{\partial}{\partial t})} \omega = \beta_\sigma^* (\xi + Gdt) \right\} \\ &= \left\{ d\beta_\sigma \left(X + F \frac{\partial}{\partial t} \right) + (\xi + Gdt) \mid \beta_\sigma^* (\xi) = e^t (i_X d\eta + F\eta), G = -e^t \eta(X) \right\}. \end{aligned}$$

Therefore, one can write $L_{M \times \mathbb{R}} \equiv \tilde{L}_M$, where L_M is the Dirac-Jacobi structure given by

$$L_M = \left\{ ((d\beta)(X), F) + (\xi, G) \mid \beta^* (\xi) = i_X d\eta + F\eta, G = -\eta(X) \right\}.$$

By construction, $\beta : G \rightarrow M$ is a Dirac-Jacobi map. There follows the result. ■

5 Examples

In this section, we will give some examples of Dirac-Jacobi structures and describe their corresponding precontact groupoids.

5.1 Precontact structures

A **precontact structure** on a manifold M is just a 1-form θ on M . A precontact structure θ on M induces a Dirac-Jacobi structure L_θ whose space of smooth sections is

$$\Gamma(L_\theta) = \{(X, f) + (i_X d\theta + f\theta, -i_X \theta) \mid (X, f) \in \mathfrak{X}(M) \times C^\infty(M)\}.$$

We observe that the Lie algebroids L_θ and $(TM \times \mathbb{R}, [\cdot, \cdot], \pi)$ are isomorphic, where $\pi : TM \times \mathbb{R} \rightarrow TM$ is the canonical projection over the first factor and $[\cdot, \cdot]$ is given by

$$[(X, f), (Y, g)] = ([X, Y], X(g) - Y(f)),$$

for $(X, f), (Y, g) \in \mathfrak{X}(M) \times C^\infty(M)$. Moreover, under the isomorphism between L_θ and $TM \times \mathbb{R}$, the 1-cocycle ϕ is the pair $(0, 1) \in \Omega^1(M) \times C^\infty(M)$.

On the other hand, consider the product $G = M \times M \times \mathbb{R}$ of the pair groupoid with $\mathbb{R}$. The function $\sigma : G \rightarrow \mathbb{R}, (x, y, t) \mapsto t$, is trivially multiplicative. In addition, if θ is a 1-form on M then one can define the 1-form η on G given by

$$\eta = \pi_1^* \theta - e^\sigma \pi_2^* \theta,$$

where $\pi_i, i \in \{1, 2\}$, is the projection on the i -th component. Then (G, η, σ) is a precontact groupoid and, moreover, the corresponding Dirac-Jacobi structure on M is just L_θ .

5.2 Dirac structures

Let L_0 be a vector subbundle of $TM \oplus T^*M$ and consider the vector subbundle L of $\mathcal{E}^1(M)$ whose sections are

$$\Gamma(L) = \{(X, 0) + (\alpha, f) \mid X + \alpha \in \Gamma(L_0), f \in C^\infty(M)\}.$$

Then, L_0 is a Dirac structure on M if and only if L is a Dirac-Jacobi structure.

If we denote by L_0 (resp., L) the Lie algebroid associated with the Dirac structure (resp., the Dirac-Jacobi structure), then we have that $L \equiv L_0 \times \mathbb{R}$. Moreover, a direct computation shows that the bracket on $\Gamma(L)$ is given by

$$[(\mathcal{X}_1, f_1), (\mathcal{X}_2, f_2)]_L = \left([[\mathcal{X}_1, \mathcal{X}_2]]_{L_0}, \rho_{L_0}(\mathcal{X}_1)(f_2) - \rho_{L_0}(\mathcal{X}_2)(f_1) + \Omega_{L_0}(\mathcal{X}_1, \mathcal{X}_2) \right),$$

for $(\mathcal{X}_1, f_1), (\mathcal{X}_2, f_2) \in \Gamma(L)$, and where $([\cdot, \cdot]_{L_0}, \rho_{L_0})$ (resp., $([\cdot, \cdot]_L, \rho_L)$) denotes the Lie algebroid structure on L_0 (resp., L) and $\Omega_{L_0} \in \Gamma(\wedge^2 L_0^*)$ is the closed 2-section given by

$$\Omega_{L_0}(\mathcal{X}_1, \mathcal{X}_2) = \frac{1}{2} \left(\xi_1(X_2) - \xi_2(X_1) \right), \text{ for } \mathcal{X}_i = X_i + \xi_i \in \Gamma(L_0).$$

Therefore, we have that the Lie algebroid structure on L is just the central extension of the Lie algebroid L_0 by the closed 2-section Ω_{L_0} . On the other hand, from Equation (2), one

sees that the 1-cocycle ϕ identically vanishes. If L_0 as well as L are integrable and (G, ω) is the presymplectic groupoid associated with L_0 then one obtains a **prequantization** of (G, ω) , that is, a central extension of Lie groupoids

$$M \times S^1 \rightarrow \tilde{G} \rightarrow G,$$

and a multiplicative 1-form $\eta \in \Omega^1(\tilde{G})$ ($m^*\eta = pr_1^*\eta + pr_2^*\eta$) which is a connection 1-form for the principal S^1 -bundle $\pi : \tilde{G} \rightarrow G$ and which satisfies $d\eta = \pi^*\omega$ (see [3]).

5.3 Jacobi manifolds

Let (Λ, E) be Jacobi manifold and $L_{(\Lambda, E)}$ the corresponding Dirac-Jacobi structure

$$L_{(\Lambda, E)} = \{(\Lambda^\sharp(\alpha_x) + \lambda E_x, -\alpha_x(E_x)) + (\alpha_x, \lambda) \mid (\alpha_x, \lambda) \in T_x^*M \times \mathbb{R}, x \in M\}.$$

In this case, the corresponding Dirac structure on $M \times \mathbb{R}$ is the one coming from the Poissonization of the Jacobi structure (Λ, E) , i.e. $\Pi = e^{-t}(\Lambda + \frac{\partial}{\partial t} \wedge E)$. Thus, the presymplectic groupoid is a honest symplectic groupoid (see [2]), and therefore, the precontact groupoid structure integrating $L_{(\Lambda, E)}$ is a **contact groupoid**, i.e., the 1-form defines a contact structure on G . This result was first proved in [6] (see also, [4, 10]). **Acknowledgments:**

D. Iglesias wishes to thank the Spanish Ministry of Education and Culture and Fulbright program for a MECD/Fulbright postdoctoral grant. A. Wade would like to thank the MSRI for hospitality while this paper was being prepared.

References

- [1] H. Bursztyn, M. Crainic: Dirac structures, momentum maps and quasi-Poisson manifolds, *The breadth of symplectic and Poisson geometry* 1-40, Progr. Math. **232**, Birkhäuser Boston, 2005.
- [2] H. Bursztyn, M. Crainic, A. Weinstein and C. Zhu: Integration of twisted Dirac brackets, *Duke Math. J.* **123** (2004) 549-607.
- [3] M. Crainic: Prequantization and Lie brackets, *preprint (2004)* math.DG/0403269.
- [4] M. Crainic and C. Zhu: Integrability of Jacobi structures, *preprint (2004)* math.DG/ 0403268.
- [5] T.J. Courant: Dirac manifolds, *Trans. A.M.S.* **319** (1990) 631-661.
- [6] P. Dazord: Sur l'intégration des algèbres de Lie locales et la préquantification, *Bull. Sci. Math.* **121** (1997) 423-462.

- [7] J. Grabowski and G. Marmo: The graded Jacobi algebras and (co)homology, *J. Phys. A: Math. Gen.* **36** (2003) 161-181.
- [8] D. Iglesias and J.C. Marrero: Lie algebroid foliations and $\mathcal{E}^1(M)$ -Dirac structures, *J. Phys. A: Math. Gen.* **35** (2002) 4085-4104.
- [9] D. Iglesias and A. Wade: Contact manifolds and generalized complex structures, *J. Geom. Phys.* **53** (2005) 249-258.
- [10] Y. Kerbrat and Z. Souici-Benhammadi: Variétés de Jacobi et groupoïdes de contact, *C. R. Acad. Sci. Paris* **317** Sér. I (1993) 81-86.
- [11] A. Wade: Conformal Dirac Structures, *Lett. Math. Phys.*, **53** (2000) 331-348.
- [12] A. Wade: Locally conformal Dirac structures and infinitesimal automorphisms, *Commun. Math. Phys.* **246** (2004) 295-310.
- [13] A. Weinstein and M. Zambon: Variations on Prequantization, *preprint (2005)* math.SG/0412502.
- [14] M. Zambon and C. Zhu: On the geometry of prequantization spaces, *paper in preparation (2005)*.